《生物科学杂志(ISSN 2958-4035)》
Journal of Biological Sciences (ISSN 2958-4035)

ChengZhu Science™

Serial in November, 2022

ChengZhu Science Research Center Limited

(Hong Kong Registration NO. 80561228)

Journal article DOI is registered via JiangXi Province ChengZhu Environmental Engineering Company Ltd, with Crossref membership (DOI prefix 10.58473)

出版人： 刘焕 香江出版社有限公司

Publisher: Liu Huan, Xiangjiang Publishing Company Ltd.

ISSN 2958-4035



Copyrights Statements

Copying and Transferring is Forbidden!

版权申明

禁止复制、转载！

All the intellectual property (mainly including the original academic knowledges and brand logo) are prohibited to copy or transfer into other publications or websites. To cite this article, only short quote is acceptable, but copying or transferring any substantial part of this article is NOT allowed (Defined in <Copyright Ordinance> in Hong Kong). The original academic knowledge is the substantial part of an article as academic journal. For learning purpose, it is allowed to read our website in online video class only. This journal is published by Hong Kong Publisher, and the copyrights is regulated and protected by <Copyright Ordinance> in Hong Kong, China. This PDF document is accessible to public only through Hong Kong domain websites (natural-foundation-science.org), and its printed version is the formally published journal. Without permission, it is NOT allowed to print, issue and sale.

所有形式知识产权（主要包括原创型学术知识和品牌标识）禁止复制、转载到其他出版物和网站。如果需要引用这篇论文，仅仅允许简短引述，但是禁止复制、转载这篇论文中任何实质性部分（香港《版权条例》中定义）。作为学术杂志，这篇论文中的原创型学术知识即为作品的实质性部分。仅仅允许以学习为目的在线视频课堂阅读本公司网站。本杂志由香港出版社出版，其版权受中国香港《版权条例》监管和保护。此PDF文档仅仅通过香港主机网站向公众公开 (natural-foundation-science.org)，并且其印刷版本杂志为正式出版物。未经许可，不得印刷、发行、销售。

Article 8: Gene Therapy and Gene Modification in Nature /基因治疗与自然界基因修改

Author: Liu Huan (1983-), Master of Science (First Class Honours, 2009), The University of Auckland.

DOI: https://doi.org/10.58473/JBS0016

Formally published on 30/11/2022; Latest revised on 26/05/2026.

Abstract

The article designs the research protocol to assess the self-repair function of genome, and potential for gene therapy and gene modification, induced by bio-signal simulation. Substantial discussion of gene therapy theory in nature is presented, rather than the 'clone' one.

Key words: Gene Self-repair, Gene Therapy, Gene Modification, Bio-signals Simulation, Biophysics.

1.The effects of electromagnetism on cells

1.1. Methods

There are three kinds of cells abstracted from the same tissue of the same genetic strain, which are cultivated in physiological saline in Lab:

Step 1. A stream of cells is cultivated under radiation condition, leading to the gene mutation of DNA molecule (sample 1) which can be detected by FISH technology; another stream of cells is cultivated in moderate electromagnetism condition leading to better immunology (sample 2) without gene mutation in DNA; the last stream of cells (sample 3) is cultivated under 'comfortable' conditions. For the simulation of moderate electromagnetism condition, cells are cultivated in electrophoresis pipe for cell electrophoresis, in which the external electric field is added. Subsequently, the electrophoresis pipe is horizontally placed under external vertical magnetism field imposed by magnetic instrument.

Step 2. The samples of even mixture between sample 1 and sample 2 (50% cell quantity for each sample) are cultivated together in moderate electromagnetism condition;

Step 3. The samples of even mixture between sample 1 and sample 3 (50% cell quantity for each sample) are cultivated together in 'comfortable' condition;

Step 4. Finally, the gene mutation rate in DNA molecules of mixed cells' sample are calculated by FISH after step 2 and step 3, respectively.

Please note: it is expected to clearly detect gene mutation after more than ten generations of cell division.

1.2. Objectives:

It is hypothesized that step 2 leads to lower gene mutation rate in mixed cells due to the assimilation of cells' 'memory' between different phenotypes discussed in previous articles [2], as compared to step 3. This is the inversion of cancerous gene mutation in DNA molecules, due to the self-repair of genome. The electric potential of different cell communities are recorded along the electrophoresis pipe in the electrophoresis instrument as discussed in another article [3], and it is expected that the cells of junior stages are more likely to be assimilated, so the advantages of this electromagnetism gradient is to separate the cell communities with different aging stages. **The electromagnetic field of cells varies during their life cycle, which is consequently a kind of time-varying electromagnetism yielding the bio-electromagnetic wave. This bio-electromagnetic wave is the bio-signal for cells to communicate each other (or for vegetation species to communicate each other).**

Please note: this method may be less effective on the cancerous cells caused by virus, due to the 'intelligence' of causal factors. The electromagnetism condition is the constant electromagnetism in this article, and the electromagnetic wave is the time-varying electromagnetism. The electromagnetic fields are obviously the constant one in this article rather than time-varying electromagnetic waves. Consequently, the method of this article is similar to the situation of 'taking rest quietly and recovery smoothly by coachers in electromagnetic field after injury,' whereas the method of application of time-varying electromagnetic waves in my another article [4], which aims to defend against invasive pathogens, is similar to the battle music violently, encouraging cells to fight.

1.3. Discussion:

Gene mutation, which occurs during DNA replication process, is triggered by the bio-signal perceived by cell. Once the bio-signal is altered, gene mutation occurs. However, the assimilation of cells results in the inversion of gene mutation during positive biophysical simulation in site due to the self-repair function of cells.

2.Gene Therapy and Gene Modification in Nature

It is further deduced that natural and mild gene mutation would occur during biophysical training process of blood cells discussed by my another article [5], which is an indicator of improved environmental adaptiveness like microbes. However, this

mild and natural gene mutation is neither like cancerous gene mutation, nor like clone cells which have been artificially inserted or deleted by other DNA sequences. As discussed in previous article, gene mutation leads to faster cell division rate. It is deduced that gene mutation caused by stronger intensity of electromagnetic waves results in faster cell division rate, so the recommended intensity of electromagnetic waves in this article is moderate and increases the blood cell activity once gene mutation is caused by this. Please keep in mind like that: gene mutation is the way for cells evolving into environmental adaptiveness in response to environmental change, and positive gene mutation can be directed in Lab, whereas negative gene mutation is prevented. To date it has been noticed that moderate gene mutation occurs after more than 10 generation cultivation using common environmental microbes in sewage water treatment, which can be detected by the above method (FISH technology is used to detect the genotype of environmental microbes in sewage water treatment plant, as a common solution to monitor the variation in environmental microbial community, and it is expected that different genotypes are evolved, which can be observed via this routine monitoring - I have observed this too).

Actually, this is easy to understand in life: let's compare two different populations: a population of Chinese who survive in city for several generations without much exposure to sunshine and the other population of Chinese who work as farmers for several generations with much exposure to sunshine. As to compare the skin color of their infants, there must be apparent different between these two populations: city populations tend to be white, and rural populations tend to be brown due to more skin melanin. Obviously, the gene-mutation-induced phenotype at skin cells' level is able to pass onto next generations as genetic materials at individual level.

Additionally, the mixed cultivation of blood cells with different genetic strains (of course, blood types must be the same) helps to improve the immunology as a whole community against invasive pathogens as well, which focus on the optimization of gene pool and enhances gene diversity in blood cell community. Please note: unlike cells of other tissues (or organs), blood cells can be the mixed cultivation from different genetic strains for 'exchange transfusion' remediation. Consequently, DNA sequencing technology is used to relate the gene variation to the specific immunology against specific pathogens, which provides basis of optimization of gene pool for cell transplantation. Further more, it is deduced that cell transplantation is more effective than the whole tissue or organ transplantation, not just because of less genetic or type matching requirement for cell transplantation, but also because cultivation of 'young' cells after transplantation would lead to less resistance against other organs. Please note: the 'young' cells are the cells with more active cell division rate caused by moderate gene mutation, and the advantages of cell transplantation is particularly important to other tissues or organs rather than just blood cells.

Further more, as discussed in previous article [5], 'the specific frequency of electromagnetic wave simulates the bio-signal regulating gene expression as a specific

isozyme family, and the specific electromagnetic wave intensity corresponds to the bio-signal regulating gene expression as a specific enzyme species within an isozyme family, which can be examined by metabolomics tests.' Consequently the specific gene locus of moderate gene mutation, expressed as specific gene trait, is caused by this biophysical training as well, which restores the specific congenital defect through gene therapy and cell transplantation. In this case, this moderate gene mutation would be understood and named as the gene variation only, because the gene mutation would be considered as higher variation by alteration of genomes.

Step 1. host cells are cultivated during simulation of electromagnetic wave conditions;

Step 2. Different frequency (or different wavelength) of electromagnetic wave are simulated, and labeled as F1, F2, ..., Fn;

Step 3. Under each simulated frequency of electromagnetic wave, different intensities of electromagnetic wave are simulated, and labeled as I1, I2, ..., and In; Then in total n×n bio-samples of host cells are cultivated under biophysical simulation;

Step 4. After sufficient reproduction process (for example, the immune cells of T lymphocytes and natural killer (NK) cells may be cultivated for approximately 10 generations or two weeks), the gene variation of each bio-samples in host cells is identified by FISH technology, recorded as different new genotypes of host cells;

Step 5. Finally, each new host cells of different genotypes is abstracted for immunology test by invasion-host interaction simulation, and then the potential gene therapy is examined. For other genetic disease, a new method of screening the specific gene sequences expressed as the specific congenital genetic defect is designed in my another article [7] to exactly target the gene sequences for gene therapy purpose through this biophysical training method.

Please note: the detection of gene variation on specific genome locus can be achieved by FISH technology, and the criterion of centromere index or the curve degree of chromosome is used to detect the specific locus of gene mutation, which becomes the unique identifier to distinguish different loci of gene mutation. However, the improved detection methods for gene variation has been designed in another structural biology article [6], and the improved simulation method of time-varying electromagnetic waves has been designed in another biophysics article [4], which can be used to replace the static intensity specified in step 3 above. It is expected that a specific life function is determined by multiple gene sequences concurrently rather than single gene sequence isolated, which means the time-varying electromagnetic wave is more reasonable than the static electromagnetic wave.

Of course, it is to further deduce the biophysics theory for gene therapy: the specific frequency of electromagnetic wave determines the gene variation in specific gene

locus qualitatively (a fragment consisted of the same and repetitive DNA sequences on chromosome), and the specific intensity of electromagnetic wave determines the 'modification' of sequence amount of these repetitive DNA sequences quantitatively on this specific gene locus. This science rhythm corresponds to the re-definition of isozyme family: the biochemistry molecules contain the same functional group in the same isozyme family, but the amount of repetitive functional groups varies between different enzyme species' molecules within an isozyme family. Obviously, specific gene locus (or loci) express as specific isozyme family in this new definition.

Please note: for the biophysics simulation of bio-signals (bio-electromagnetic waves), it is important to know the parameters about the frequency range of bio-electromagnetic waves released by human species or other biological species. Although bio-electromagnetic wave detector becomes the common equipment now, as summarized by my another article [8], the detailed parameters of bio-electromagnetic wave frequency range for each species is still limited in online published papers (I have searched this information via Google, GPT-5.5 and Claude Opus 4.6, but the information is limited, and this information may be usually used to design patent instead). Consequently, it is still necessary for researchers to measure these parameters by themselves, but it is believed that the detector of bio-electromagnetic wave is already convenient to access now.

References:

[1] 注册环保工程师专业考试复习教材(2009). 第二分册. 中国环境科学出版社. ISBN：978-7-5111-0505-9.

[2]. Liu Huan. (2021). Assimilation of The Cells' 'Memory' Between Different Phenotypes and Its Implication on Canceration. Journal of Environment and Health Science (ISSN 2314-1628), 2021(02).https://doi.org/10.58473/JBS0015

[3]. Liu Huan. (2021). Bio-magnetic field of Cell and Its Application on Separation of Blood Cell Communities along Environmental Gradient. Journal of Environment and Health Science (ISSN 2314-1628), 2021(02).https://doi.org/10.58473/JBS0011

[4]. Liu Huan. (2021). The Parameterization of Time-varying Electromagnetic Field for Biophysics Simulation. Journal of Environment and Health Science (ISSN 2314-1628), 2021(02).https://doi.org/10.58473/JBS0013

[5]. Liu Huan. (2021). Bio-signal Simulation of Electromagnetic Wave and Its Specificity on the Isozyme Expression. Journal of Environment and Health Science (ISSN 2314-1628), 2021(02).https://doi.org/10.58473/JBS0012

[6]. Liu Huan. (2022). Essay: Structural Molecular Biology. Journal of Environment and Health Science (ISSN 2314-1628), 2022(08).https://doi.org/10.58473/JBS0021

[7]. Liu Huan. (2023). Original review of specificity in the interaction between pathogen invasion and host organism. Journal of Biological Sciences (ISSN 2958-4035). 2023 (04). https://doi.org/10.58473/JBS0022

[8]. Liu Huan. (2021). The Synthesis of Biological Antibiotics and Its Application on Bio-medicine. Journal of Environment and Health Science (ISSN 2314-1628), 2021(02). https://doi.org/10.58473/JBS0014